

KEK-TH-568
 hep-ph/9804327
 April 1998

Impacts on searching for signatures of new physics from $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay[†]

Gi-Chol Cho[‡]

Theory Group, KEK, Tsukuba, Ibaraki 305, Japan

Abstract

Impacts on new physics search from B^0 - $\bar{B}^0$, K^0 - $\bar{K}^0$ mixings and the rare decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ are discussed. We show that, in a certain class of new physics models, the extra contributions to those processes can be parametrized by its ratio to the standard model (SM) contribution with the common CKM factors. We introduce two ratios to measure the new physics contributions, R_1 for x_d and ϵ_K parameters, and R_2 for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ decay. Then, the experimentally allowed region for the new physics contributions can be given in terms of R_1 , R_2 and the CP violating phase δ of the CKM matrix. We find constraints on R_1 and $\cos \delta$ by taking account of current experimental data and theoretical uncertainties on B^0 - $\bar{B}^0$ and K^0 - $\bar{K}^0$ mixings. We also study impacts of future improved measurements on $(R_1, R_2, \cos \delta)$ basis. As typical examples of new physics models, we examine contributions to those processes in the minimal supersymmetric SM and the two Higgs doublet model.

[†]Talk given at the workshop on “Fermion Mass and CP Violation”, Hiroshima, Japan, 5-6 March 1998.

[‡]Research Fellow of the Japan Society for the Promotion of Science

1 Introduction

Processes mediated by flavor changing neutral current (FCNC) have been considered as good probes of physics beyond the standard model (SM). By using the experimentally well measured processes, it is expected to obtain an indirect evidence or constraints on new physics models. An existence of new physics may arise as violation of the unitarity of the Cabbibo-Kobayashi-Maskawa (CKM) matrix. Such signatures of new physics will be explored through the determination of the unitarity triangle at B-factories at KEK and SLAC in the near future.

Typical FCNC processes which have been often used to study the new physics contributions are $B^0\text{-}\overline{B}^0$ and $K^0\text{-}\overline{K}^0$ mixings. Parameters x_d in $B^0\text{-}\overline{B}^0$ mixing and ϵ_K in $K^0\text{-}\overline{K}^0$ mixing are dominated by the short distance physics and have been calculated in the SM and many new physics models. Experimentally, both parameters have been measured as [1]

$$x_d = 0.73 \pm 0.05, \quad (1.1a)$$

$$|\epsilon_K| = (2.23 \pm 0.013) \times 10^{-3}. \quad (1.1b)$$

On the other hand, there are large theoretical uncertainties on both parameters which come from the evaluation of the hadronic matrix elements of those processes. They are parametrized in terms of decay constants and bag-parameters of B or K mesons. Thus, the loss of information on the CKM matrix elements or the new physics contributions from those processes is not avoidable in the level of these uncertainties.

The rare decay $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ is one of the most promising processes to extract clean informations about the CKM matrix elements [2] because the decay rate of this process has small theoretical uncertainties. The reason can be summarized as:

- The process is dominated by the short-distance physics. The long-distance contributions have been estimated as 10^{-3} smaller than the short-distance contributions [3].
- The hadronic matrix element of the decay rate can be evaluated by using that of $K^+ \rightarrow \pi^0 e^+ \nu_e$ process which is accurately measured.
- The short distance contributions in the SM have been calculated in the next-to-leading order (NLO) level (for a review, see [2]).

With these attractive points, the first observation of an event consistent with this decay process which was reported by E787 collaboration [4]

$$\text{Br}(K^+ \rightarrow \pi^+ \nu \overline{\nu}) = 4.2_{-3.5}^{+9.7} \times 10^{-10}, \quad (1.2)$$

motivates us to examine the implication of the above estimate of the branching fraction and of its improvement in the near future.

In this report, we discuss impacts on the search for a new physics signal from the above FCNC processes – $B^0\text{-}\overline{B}^0$, $K^0\text{-}\overline{K}^0$ mixings and $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ decay. We focus on a class of new physics models which satisfy the following three conditions:

- (i) FCNC in the new physics sector is described by the $V - A$ type operator.
- (ii) The flavor mixing in the new physics sector is governed by the SM CKM matrix elements.
- (iii) The net contributions are proportional to the CKM matrix elements which are concerned with the third generation.

In the following, we will show that new physics contributions to those processes can be parametrized by two quantities, R_1 for $B^0\text{-}\overline{B}^0, K^0\text{-}\overline{K}^0$ mixings and R_2 for $K^+ \rightarrow \pi^+\nu\overline{\nu}$ decay. Both quantities are defined as the ratio of the new physics contribution to that of the SM. Constraints on the new physics contributions are summarized in terms of R_1, R_2 and $\cos\delta$, where δ is the CP violating phase of the CKM matrix in the standard parametrization [1]. Taking account of current experimental data on x_d and ϵ_K parameters in $B^0\text{-}\overline{B}^0$ and $K^0\text{-}\overline{K}^0$ mixings, and uncertainties in the hadronic parameters, we will show constraints on R_1 and $\cos\delta$. In order to see that how $K^+ \rightarrow \pi^+\nu\overline{\nu}$ decay could give impacts on new physics search, we will find constraints on R_1, R_2 and $\cos\delta$ by assuming the future improvement of the $\text{Br}(K^+ \rightarrow \pi^+\nu\overline{\nu})$ measurements. As examples of new physics models which naturally satisfy the above three conditions, we will find the consequences of the minimal supersymmetric standard model (MSSM) [5] and the two Higgs doublet model (THDM) [6].

2 New physics contributions to the FCNC processes in the B and K meson systems

The effective Lagrangian for the $K^+ \rightarrow \pi^+\nu\overline{\nu}$ process in the SM is given by [7]:

$$\mathcal{L}_{eff}^{K^+} = \frac{G_F}{\sqrt{2}} \frac{2\alpha(m_Z)}{\pi} \frac{1}{\sin^2\theta_W} \overline{\nu}_\ell \gamma^\mu P_L \nu_\ell \overline{s} \gamma_\mu P_L d \sum_{i=2,3} V_{i2}^* V_{i1} \eta_i D_W(i), \quad (2.1)$$

where i and ℓ are the generation indices for the up-type quarks and leptons, respectively. The CKM matrix element is given by V_{ij} and the projection operator P_L is defined as $P_L \equiv (1 - \gamma_5)/2$. The loop function for the i -th generation quark is denoted by $D_W(i)$ and its explicit form can be found in [7]. The QCD correction factor for the top-quark exchange has been estimated as $\eta_3 = 0.985$ for $170 \text{ GeV} \leq m_t \leq 190 \text{ GeV}$ [8]. The QCD correction factor for the charm-quark exchange with its loop function is numerically given as $\eta_2 D_W(2) = \lambda^4 \times (0.40 \pm 0.06)$ [9] where $\lambda \equiv |V_{12}|$. The error is due to uncertainties in the charm quark mass and higher order QCD corrections. Then, summing up the three generations of neutrino, the branching ratio is expressed as [10]

$$\text{Br}(K^+ \rightarrow \pi^+\nu\overline{\nu}) = 1.57 \times 10^{-4} \left| V_{32}^* V_{31} \eta_3 D_W(3) + V_{22}^* V_{21} \eta_2 D_W(2) \right|^2. \quad (2.2)$$

With the above estimates for the loop functions and the QCD correction factors, the branching ratio is predicted to be [11]

$$\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{SM}} = (9.1 \pm 3.8) \times 10^{-11} \quad (2.3)$$

in the SM, where the error is dominated by the uncertainties of the CKM matrix elements.

The effective Lagrangian of the B^0 - $\bar{B}^0$ mixing in the SM is expressed by

$$\mathcal{L}_{eff}^{\Delta B=2} = \frac{G_F^2 M_W^2}{4\pi^2} \bar{d}\gamma^\mu P_L b \bar{d}\gamma_\mu P_L b \sum_{i,j=2,3} V_{i1}^* V_{i3} V_{j1}^* V_{j3} F_V^W(i, j). \quad (2.4)$$

Likewise, $\mathcal{L}_{eff}^{\Delta S=2}$ for the K^0 - $\bar{K}^0$ mixing is obtained by replacing V_{i3} with V_{i2} , and the b -quark operators with the s -quark ones, respectively. The explicit form of the loop function $F_V^W(i, j)$ is given in [7]. The B -meson mixing parameter x_d is defined by $x_d \equiv \Delta M_B / \Gamma_B$, where ΔM_B and Γ_B correspond to the B -meson mass difference and the average width of the mass eigenstates, respectively. The mass difference is induced by the above $\Delta B = 2$ operator (2.4) and we can express the mixing parameter x_d in the SM as

$$x_d = \frac{G_F^2}{6\pi^2} M_W^2 \frac{M_B}{\Gamma_B} f_B^2 B_B |V_{31}^* V_{33}|^2 \eta_B |F_V^W(3, 3)|, \quad (2.5)$$

where f_B , B_B and η_B denote the decay constant of B^0 -meson, the bag parameter of B^0 - $\bar{B}^0$ mixing and the short-distance QCD correction factor, respectively.

The CP-violating parameter ϵ_K in the K^0 - $\bar{K}^0$ system is given by the imaginary part of the same box diagram of the B^0 - $\bar{B}^0$ transition besides the external quark lines. We can express the ϵ_K parameter in the SM as

$$\begin{aligned} \epsilon_K = & -e^{i\pi/4} \frac{G_F^2}{12\sqrt{2}\pi^2} M_W^2 \frac{M_K}{\Delta M_K} f_K^2 B_K \text{Im} \left\{ (V_{31}^* V_{32})^2 \eta_{K33} F_V^W(3, 3) \right. \\ & \left. + (V_{21}^* V_{22})^2 \eta_{K22} F_V^W(2, 2) + 2(V_{31}^* V_{32} V_{21}^* V_{22}) \eta_{K32} F_V^W(3, 2) \right\}, \end{aligned} \quad (2.6)$$

where f_K , B_K and η_{Kij} represent the decay constant, the bag parameter and the QCD correction factors, respectively.

In theoretical estimation of these quantities, non-negligible uncertainties come from the evaluations of the QCD correction factors and the hadronic matrix elements. In our analysis, we adopt the following values:

$$\eta_B = 0.55 \pm 0.01 [12], \quad \sqrt{B_B} f_B = (220 \pm 40) \text{ MeV} [13], \quad (2.7)$$

for the x_d parameter, and

$$\left. \begin{aligned} \eta_{K33} &= 0.57 \pm 0.01 \\ \eta_{K22} &= 1.38 \pm 0.20 \\ \eta_{K32} &= 0.47 \pm 0.04 \end{aligned} \right\} [12, 14], \quad B_K = 0.75 \pm 0.15 [11]. \quad (2.8)$$

for the ϵ_K parameter.

Next, we consider the new physics contributions to these quantities, $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ (2.2), x_d (2.5), and ϵ_K (2.6). In a class of new physics models which satisfy our three conditions, the effective Lagrangians can be obtained by replacing $D_W(i)$ with $D^{\text{new}}(i)$ in (2.1), and $F_V^W(i, j)$ with $F_V^{\text{new}}(i, j)$ in (2.5) and (2.6). Then, the effective Lagrangians of these processes in the new physics sector should have the following forms;

$$\mathcal{L}_{\text{new}}^{K^+} = \frac{G_F}{\sqrt{2}} \frac{2\alpha(m_Z)}{\pi} \frac{1}{\sin^2 \theta_W} \bar{\nu} \gamma^\mu P_L \nu \ V_{32}^* V_{31} \ \bar{s} \gamma_\mu P_L d \ A^{\text{new}}, \quad (2.9a)$$

$$\mathcal{L}_{\text{new}}^{\Delta B=2} = \frac{G_F^2 M_W^2}{4\pi^2} \bar{d} \gamma^\mu P_L b \ \bar{d} \gamma_\mu P_L b \ (V_{31}^* V_{33})^2 \ B^{\text{new}}, \quad (2.9b)$$

$$\mathcal{L}_{\text{new}}^{\Delta S=2} = \frac{G_F^2 M_W^2}{4\pi^2} \bar{d} \gamma^\mu P_L s \ \bar{d} \gamma_\mu P_L s \ (V_{31}^* V_{32})^2 \ B^{\text{new}}. \quad (2.9c)$$

It should be noticed that the new physics contributions to the $\Delta B = 2$ (2.9b) and the $\Delta S = 2$ (2.9c) processes are expressed by the same quantity B^{new} .

There are two cases in which the effective Lagrangians can be given by the above forms. First, if the contributions from the first two generations do not differ much, *i.e.*,

$$D^{\text{new}}(2) \approx D^{\text{new}}(1), \quad (2.10a)$$

$$F_V^{\text{new}}(i, 1) \approx F_V^{\text{new}}(i, 2), \quad (2.10b)$$

the net contributions from the new physics are written by using the unitarity of the CKM matrix as;

$$\sum_i V_{i2}^* V_{i1} D^{\text{new}}(i) \approx V_{32}^* V_{31} \{D^{\text{new}}(3) - D^{\text{new}}(1)\}, \quad (2.11a)$$

$$\begin{aligned} \sum_{i,j} V_{i1}^* V_{ik} V_{j1}^* V_{jk} F_V^{\text{new}}(i, j) &\approx (V_{31}^* V_{3k})^2 \{F_V^{\text{new}}(3, 3) + F_V^{\text{new}}(1, 1) \\ &\quad - F_V^{\text{new}}(3, 1) - F_V^{\text{new}}(1, 3)\}, \end{aligned} \quad (2.11b)$$

for $k = 2, 3$. We can now define the parameters A^{new} and B^{new} as

$$A^{\text{new}} \equiv D^{\text{new}}(3) - D^{\text{new}}(1), \quad (2.12a)$$

$$B^{\text{new}} \equiv F_V^{\text{new}}(3, 3) + F_V^{\text{new}}(1, 1) - F_V^{\text{new}}(3, 1) - F_V^{\text{new}}(1, 3). \quad (2.12b)$$

Second, if the contributions from both the first two generations are negligible as compared with those of the 3rd generation, *i.e.*,

$$D^{\text{new}}(3) \gg D^{\text{new}}(1), D^{\text{new}}(2), \quad (2.13a)$$

$$F_V^{\text{new}}(3, 3) \gg F_V^{\text{new}}(1, j), F_V^{\text{new}}(2, j), F_V^{\text{new}}(3, 1), F_V^{\text{new}}(3, 2), \quad (2.13b)$$

the parameters A^{new} and B^{new} become

$$A^{\text{new}} = D^{\text{new}}(3), \quad (2.14a)$$

$$B^{\text{new}} = F_V^{\text{new}}(3, 3). \quad (2.14b)$$

Now, the effects of the new physics contributions to these processes can be evaluated by the following ratios [15]¹

$$R_1 = \frac{F_V^W(3, 3) + B^{\text{new}}}{F_V^W(3, 3)}, \quad (2.15a)$$

$$R_2 = \frac{D_W(3) + A^{\text{new}}}{D_W(3)}. \quad (2.15b)$$

Once a model of new physics is specified, we can quantitatively estimate its effect in terms of R_1 and R_2 . Both parameters converge to unity as the new physics contributions are negligible,

$$R_1, R_2 \longrightarrow 1 \quad \text{for} \quad A^{\text{new}}, B^{\text{new}} \longrightarrow 0. \quad (2.16)$$

Because constraint on R_2 is obtained from $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, it can be a negative quantity if the extra contributions destructively interfere with that of the SM. In the following, we consider the cases where the net contributions from the new physics sector do not exceed those of the SM: $A^{\text{new}} < |D_W(3)|$ and $B^{\text{new}} < |F_V^W(3, 3)|$. Then, we study constraints on R_1 and R_2 from experimental data in the range of $0 < R_1, R_2 < 2$.

3 Constraints on new physics contributions to FCNC processes

Sizable new physics effects to x_d, ϵ_K and $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ can be detected as deviations of R_1 and R_2 from unity. In practice, experimentally measurable quantities are products of R_1 or R_2 by the CKM matrix elements. In the standard parametrization of the CKM matrix, the uncertainty in the CP-violating phase δ dominates that of the CKM matrix elements [1]. Hence, together with R_1 and R_2 , we allow $\cos \delta$ to be fitted by the measurements of x_d, ϵ_K and $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$. By this reason, constraints on R_1 and R_2 are correlated through $\cos \delta$.

We perform the χ^2 -fit for two parameters R_1 and $\cos \delta$ by using experimental data of x_d and ϵ_K . In the fit, we take into account of the theoretical uncertainties which are given in (2.7), (2.8) and

$$\left. \begin{aligned} |V_{12}| &= 0.2205 \\ |V_{23}| &= 0.041 \pm 0.003 \\ |V_{13}/V_{23}| &= 0.08 \pm 0.02 \end{aligned} \right\} [1], \quad m_t = 175.6 \pm 5.5 \text{ GeV} [17], \quad (3.1)$$

where the error of $|V_{12}|$ can be safely neglected. We find

$$\left. \begin{aligned} \cos \delta &= 0.36 \pm 0.83 \\ R_1 &= 0.93 \pm 0.75 \end{aligned} \right\} \rho_{\text{corr}} = 0.90. \quad (3.2)$$

¹ Similar parametrization was used in [16]. In the article, the both ratios were defined as complex parameters. In that case, there are two additional parameters – two complex phases of these ratios.




Figure 1: The 1- σ (39% CL) allowed region from the experimental results of the $B^0\text{-}\overline{B}^0$, $K^0\text{-}\overline{K}^0$ mixings. The range between the two solid lines is the allowed region of $\cos \delta$ in the SM.

Because of the strong positive correlation between the errors, only the following combination is effectively constrained;

$$R_1 = 0.61 + 0.89 \cos \delta \pm 0.33. \quad (3.3)$$

We show the 1- σ (39%) allowed region of $\cos \delta$ and R_1 in Fig. 1. In the figure, there is small region which corresponds to $1 \leq \cos \delta$ where the flavor mixing does not obey the CKM mechanism. The range of $\cos \delta$ along the $R_1 = 1$ line is the allowed region of $\cos \delta$ in the SM: $0.08 \lesssim \cos \delta \lesssim 0.78$. We can read off from Fig. 1 that the current experimental data of x_d and ϵ_K parameters constrain the new physics contributions within $0.18 \lesssim R_1 \lesssim 1.68$.

Next we examine the constraint on R_2 . Although the recent observation of one candidate event is unsuitable to include in the actual fit, we can expect that the data will be improved in the near future. In the following, we adopt the central value of the SM prediction as the mean value of $\text{Br}(K^+ \rightarrow \pi^+ \nu \overline{\nu})$ and study consequences of improved measurements. With several more events, the branching fraction can be measured as $\text{Br}(K^+ \rightarrow \pi^+ \nu \overline{\nu}) = (0.9 \pm 0.4) \times 10^{-10}$. Then the combined result with x_d and ϵ_K parameters can be found as

$$\left. \begin{array}{l} \cos \delta = 0.36 \pm 0.83 \\ R_1 = 0.93 \pm 0.75 \\ R_2 = 1.14 \pm 0.53 \end{array} \right\} \rho_{\text{corr}} = \begin{pmatrix} 1 & 0.90 & 0.68 \\ & 1 & 0.62 \\ & & 1 \end{pmatrix}. \quad (3.4)$$




Figure 2: The $1\text{-}\sigma$ allowed regions of R_1, R_2 parameters. Three contours are corresponding to $\cos \delta = 0.36$ (solid line), $\cos \delta = -0.47$ (dotted line) and $\cos \delta = 1.19$ (dashed line), respectively.

In Fig. 2, the results are shown on the R_1 - R_2 plane for three values of $\cos \delta$; $\cos \delta = 0.36$ (mean value), -0.47 (mean value -1σ) and 1.19 (mean value $+1\sigma$). Using this result, we can discuss about constraints on the new physics contributions to these processes on the R_1 - R_2 plane for a given value of $\cos \delta$.

4 Predictions on R_1, R_2 in the MSSM and the THDM

Here, we find predictions on R_1, R_2 in the MSSM and the THDM. The previous studies on those processes in both models can be found in [18, 19, 20, 21] for B^0 - $\overline{B}^0, K^0$ - $\overline{K}^0$ mixings, and [16, 22, 23] for $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ process.

In the MSSM based on $N = 1$ supergravity [5], there are several extra particles. Then, interactions among them could be new sources of FCNC processes. It is known that chargino $-t$ -squark exchange and charged Higgs- t -quark exchange processes give the leading contributions to FCNC processes for B and K meson systems. For the chargino contribution, effects from squarks in the first two generations are canceled each other because degeneracy among their masses holds in good approximation. The interactions among the charged Higgs boson and the up-type quarks are the same with those of the type II-THDM [6]. The charged Higgs boson interacts with the up-type quarks through the Yukawa interactions which are proportional to the corresponding quark masses. As a result, the charged




Figure 3: The MSSM (left) and THDM (right) contributions to R_1 and R_2 parameters for $\tan \beta = 2$. The $1\text{-}\sigma$ allowed region of R_1 and R_2 parameters for $\cos \delta = 0.36$ is also shown.

Higgs contributions to the FCNC processes are dominated by its interaction with the top-quark.

The magnitudes of both the chargino and the charged Higgs contributions are proportional to $1/\tan \beta$, where $\tan \beta$ is the ratio of the vacuum expectation values of two Higgs fields. The effective Lagrangians for both contributions are described by $V - A$ and $S + P$ operators. The latter can be negligible for small $\tan \beta$. Furthermore, contributions from other sources in the MSSM do not give sizable effects to the FCNC processes for $\tan \beta \lesssim 10$ [16, 20]. Hence we examine both models in the region $\tan \beta \lesssim 10$.

The expressions for R_1 in the MSSM and the THDM can be found in [19]. The MSSM contribution to the decay process $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ is expressed by using D^{new} as follows

$$D^{\text{new}}(i) = \sum_{m,n,k,\alpha,\beta} D_C(i, m, n; \ell, k; \alpha, \beta) + D_H(i, \ell), \quad (4.1)$$

where $D_C(i, m, n; \ell, k; \alpha, \beta)$ and $D_H(i, \ell)$ represent the chargino and the charged Higgs boson contributions, respectively. Their explicit forms are given in [15]. For D_C , by using the unitarity of the CKM matrix and the degeneracy of the squark masses between the first two generations, we obtain

$$V_{i2}^* V_{i1} D_C(i, m, n; \ell, k; \alpha, \beta) = V_{32}^* V_{31} \left\{ D_C(3, m, n; \ell, k; \alpha, \beta) - D_C(1, m, n; \ell, k; \alpha, \beta) \right\}, \quad (4.2)$$

and the chargino contribution $A^{\text{new}} \equiv A_C$ is given by

$$A_C \equiv \sum_{m,n,k,\alpha,\beta} \left\{ D_C(3, m, n; \ell, k; \alpha, \beta) - D_C(1, m, n; \ell, k; \alpha, \beta) \right\}. \quad (4.3)$$

For D_H , due to the smallness of the Yukawa couplings for light quarks, we can write the charged Higgs contribution as

$$A_H \equiv D_H(3, \ell). \quad (4.4)$$

From (4.3) and (4.4), R_2 in the MSSM is defined as

$$R_2 \equiv \frac{D_W(3) + A_C + A_H}{D_W(3)}. \quad (4.5)$$

On the other hand, the THDM contribution to R_2 is given by setting $D_C = 0$ in (4.5):

$$R_2 \equiv \frac{D_W(3) + A_H}{D_W(3)}. \quad (4.6)$$

Let us proceed numerical study. In order to reduce the number of input parameters in the MSSM, we express the soft SUSY breaking scalar masses in the sfermion sector by a common mass parameter m_0 . Also taking the scalar trilinear coupling A_f for sfermion $\tilde{f}$ as $A_f = m_0$, the MSSM contributions can be evaluated by using four parameters, $m_0, \tan \beta$, the higgsino mass term μ and the SU(2) gaugino mass term m_2 . In our study, these parameters are taken to be real. In Fig. 3, we show the MSSM and THDM contributions to R_1, R_2 parameters with the constraints on these parameters for $\cos \delta = 0.36$. The numerical study was performed in the range of $100 \text{ GeV} < m_0 < 1 \text{ TeV}$, $|\mu| < 200 \text{ GeV}$ and $m_2 = 200 \text{ GeV}$ for $\tan \beta = 2$. We fixed the charged Higgs boson mass at $m_H = 200 \text{ GeV}$ in the MSSM prediction. This is the reason why the MSSM contributions do not converge to $R_1 = 1$ in the figure. We take into account the recent estimation of lower mass limits for lighter t -squark and lighter chargino [24]: $80 \text{ GeV} \leq m_{\tilde{t}_1}$ and $91 \text{ GeV} \leq m_{\tilde{\chi}_1}$. The MSSM contribution to R_1 interferes with that of the SM constructively [19, 20, 25]. On the other hand, the contribution to R_2 interferes with that of the SM both constructively and destructively. Contrary to the case of the MSSM, the THDM contribution constructively interferes with the SM contribution for both R_1 and R_2 . The Yukawa interaction between the top-quark and the charged Higgs boson is proportional to $1/\tan^2 \beta$. Thus constraints on the THDM contribution to these quantities are weakened together with the increase of $\tan \beta$.

5 Summary

We have studied impacts on searching for signatures of new physics beyond the SM from some FCNC processes – $B^0\text{-}\overline{B}^0, K^0\text{-}\overline{K}^0$ mixings and the rare decay $K^+ \rightarrow \pi^+ \nu \overline{\nu}$. For a certain class of models of new physics, two parameters R_1 and R_2 were introduced to estimate the new physics contributions to $B^0\text{-}\overline{B}^0, K^0\text{-}\overline{K}^0$ mixings and $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ decay, respectively. Then constraints on the new physics contributions are obtained from experimental data by using these parameters and $\cos \delta$.

Taking account of both experimental and theoretical uncertainties for the B^0 - $\overline{B}^0$ and K^0 - $\overline{K}^0$ mixings, we found current constraint on R_1 as $0.18 \lesssim R_1 \lesssim 1.68$. With the assumption that the future data of $\text{Br}(K^+ \rightarrow \pi^+ \nu \overline{\nu})$ will be close to the SM prediction, constraints on $\cos \delta$, R_1 and R_2 were found. The results were applied to the MSSM and the THDM contributions to those processes. Although there are parameter space which give roughly 50% enhancement of R_1 , contributions to R_2 are less than $\pm 10\%$. So quite precise experimental measurement of $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ is required to study constraints on the parameter space of these models.

Acknowledgment

The author would like to thank T. Morozumi and the organizing staffs of the workshop for making an opportunity to give his talk. This work is supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

References

- [1] Particle Data Group, R.M. Barnett *et al.*, Phys. Rev. **D54** (1996) 1.
- [2] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125.
- [3] J. Ellis and J.S. Hagelin, Nucl. Phys. **B217** (1983) 189;
D. Rein and L.M. Sehgal, Phys. Rev. **D39** (1989) 3325;
J.S. Hagelin and L.S. Littenberg, Prog. Part. Nucl. Phys. **23** (1989) 1;
C.Q. Gang, I.J. Hsu and Y.C. Lin, Phys. Lett. **B355** (1995) 569;
S. Fajfer, Nuov. Cim. **A110** (1997) 397.
- [4] E787 collaboration, Phys. Rev. Lett. **79** (1997) 2204.
- [5] For reviews, see, H.P. Nilles, Phys. Rep. **110** (1984) 1,
H.E. Haber and G.L. Kane, Phys. Rep. **117** (1985) 75.
- [6] See, *e.g.*, J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs Hunter's Guide*, Addison-Wesley, (1990) and references therein.
- [7] T. Inami and C.S. Lim, Prog. Theor. Phys. **65** (1981) 297; (E) 1772.
- [8] G. Buchalla and A.J. Buras, Nucl. Phys. **B400** (1993) 225.
- [9] G. Buchalla and A.J. Buras, Phys. Rev. **D54** (1996) 6782.
- [10] W.J. Marciano and Z. Parsa, Phys. Rev. **D53** (1996) R1.

- [11] A.J. Buras and R. Fleischer, hep-ph/9704376;
A.J. Buras, hep-ph/9711217.
- [12] A.J. Buras, M. Jamin and P.H. Weisz, Nucl. Phys. **B347** (1990) 491.
- [13] A. Abada, *et al.*, Nucl. Phys. **B376** (1992) 172. A. Abada, LPTHE Orsay-94/57.
- [14] S. Herrlich and U. Nierste, Nucl. Phys. **B419** (1994) 292;
S. Herrlich and U. Nierste, Phys. Rev. **D52** (1995) 6505, Nucl. Phys. **B476** (1996) 27.
- [15] G.C. Cho, to appear in Euro. Phys. Journal. **C**.
- [16] A.J. Buras, A. Romanio and L. Silvestrini, hep-ph/9712398.
- [17] CDF Collaboration, J. Lys, talk at ICHEP96, in Proc. of ICHEP96, (ed) Z. Ajduk and A.K. Wroblewski, World Scientific, (1997);
D0 Collaboration, S. Protopopescu, talk at ICHEP96, in the proceedings.
P. Tipton, talk at ICHEP96, in the proceedings.
- [18] T. Kurimoto, Phys. Rev. **D39** (1989) 3447;
S. Bertolini, F. Borzumati, A. Masiero and G. Ridolfi, Nucl. Phys. **B353** (1991) 591;
G. Couture and H. König, Z. Phys. **C69** (1996) 499.
- [19] G.C. Branco, G.C. Cho, Y. Kizukuri and N. Oshimo, Phys. Lett. **B337** (1994) 316; Nucl. Phys. **B449** (1995) 483.
- [20] T. Goto, T. Nihei and Y. Okada, Phys. Rev. **D53** (1996) 5233; (E) **D54** (1996) 5904.
- [21] L.F. Abbott, P. Sikivie and M.B. Wise, Phys. Rev. **D21** (1980) 1393.
- [22] S. Bertolini and A. Masiero, Phys. Lett. **B174** (1986) 343;
B. Mukhopadhyaya and A. Raychaudhuri, Phys. Lett. **B189** (1987) 203;
I.I. Bigi and F. Gabbiani, Nucl. Phys. **B367** (1991) 3;
G. Couture and H. König, Z. Phys. **C69** (1995) 167;
Y. Nir and M.P. Worah, hep-ph/9711215.
- [23] A.J. Buras, P. Krawczyk, M.E. Lautenbacher and C. Salazar, Nucl. Phys. **B337** (1990) 284.
- [24] P. Janot, talk given at International Europhysics Conference on High Energy Physics, Jerusalem 1997.
- [25] T. Kurimoto, Mod. Phys. Lett. **A10** (1995) 1577.